

α -Discounting Method for Multi-Criteria Decision Making (α -DMCDM)

Florentin Smarandache ¹⁹

Dedicated to Themistocles M. Rassias on the occasion of his 75th Birthday

ABSTRACT. In this paper we introduce a new procedure called α -Discounting Method for Multi- Criteria Decision Making (α -D MCDM), which is as an alternative and extension of Saaty's Analytical Hierarchy Process (AHP). It works for any number of preferences that can be transformed into a system of homogeneous linear equations. A degree of consistency (and implicitly a degree of inconsistency) of a decision-making problem are defined. α -D MCDM is generalized to a set of preferences that can be transformed into a system of linear and/or non-linear homogeneous and/or non-homogeneous equations and/or inequalities. Many consistent, weak inconsistent, and strong inconsistent examples are given.

1 INTRODUCTION

α -Discounting Method for Multi-Criteria Decision Making (α -D MCDM) is an alternative and extension of Saaty's Analytical Hierarchy Process (AHP)

- see [1-11] for more information on AHP since it is not the main subject of this paper. It works not only for preferences that are pairwise comparisons of criteria as AHP does, but for preferences of any n -wise (with $n \geq 2$) comparisons of criteria that can be expressed as linear homogeneous equations.

The general idea of α -D MCDM is to assign non-null positive parameters $\alpha_1, \alpha_2, \dots, \alpha_p$ to the coefficients in the right-hand side of each preference that

¹⁹Received: 23.08.2025

2010 *Mathematics Subject Classification.* 20F10.

Key words and phrases. Multi-Criteria Decision Making (MCDM), Analytical Hierarchy Process (AHP), α - Discounting Method, Fairness Principle, parameterize, pairwise comparison, n -wise comparison, consistent MCDM problem, weak or strong inconsistent MCDM problem.

diminish or increase them in order to transform the above linear homogeneous system of equations which has only the null-solution, into a system having a particular non-null solution.

After finding the general solution of this system, the principles used to assign particular values to all parameters α 's is the second important part of α -D, yet to be deeper investigated in the future.

In the current paper we herein propose the Fairness Principle, i.e. each coefficient should be discounted with the same percentage (we think this is fair: not making any favouritism or unfairness to any coefficient), but the reader can propose other principles.

For consistent decision-making problems with pairwise comparisons, α -Discounting Method together with the Fairness Principle give the same result as AHP.

But for weak inconsistent decision-making problem, α -Discounting together with the Fairness Principle give a different result from AHP.

α -Discounting/Fairness-Principle together give a justifiable result for strong inconsistent decision-making problems with two preferences and two criteria; but for more than two preferences with more than two criteria and the Fairness Principle has to be replaced by another principle of assigning numerical values to all parameters α 's.

Since Saaty's AHP is not the topic of this paper, we only recall the main steps of applying this method, so the results of α -D MCDM and of AHP could be compared.

AHP works only for pairwise comparisons of criteria, from which a square Preference Matrix, A (of size $n \times n$), is built. Then one computes the maximum eigenvalue λ_{max} of A and its corresponding eigenvector.

If λ_{max} is equal to the size of the square matrix, then the decision-making problem is consistent, and its corresponding normalized eigenvector (Perron-Frobenius vector) is the priority vector.

If λ_{max} is strictly greater than the size of the square matrix, then the

decision-making problem is inconsistent. One raise to the second power matrix A, and again the resulted matrix is raised to the second power, etc. obtaining the sequence of matrices $A^2, A^4, A^8, \dots$, etc.

In each case, one computes the maximum eigenvalue and its associated normalized eigenvector, until the difference between two successive normalized eigenvectors is smaller than a given threshold. The last such normalized eigenvector will be the priority vector. Saaty defined the Consistency Index as:

$$CI(A) = \frac{\lambda_{max}(A) - n}{n - 1}$$

where n is the size of the square matrix A.

2 α -Discounting Method for Multi-Criteria Decision Making (α -D MCDM).

2.1. Description of α -D MCDM.

The general idea of this paper is to discount the coefficients of an inconsistent problem to some percentages in order to transform it into a consistent problem.

Let the Set of Criteria be $= \{C_1, C_2, \dots, C_n\}$, with $n \geq 2$, and the Set of Preferences be $P = \{P_1, P_2, \dots, P_m\}$, with $m \geq 1$.

Each preference P_i is a linear homogeneous equation of the above criteria $C_1, C_2, \dots, C_n$:

$$P_i = f(C_1, C_2, \dots, C_n)$$

We need to construct a basic belief assignment (bba):

$$m : C \rightarrow [0, 1]$$

such that $m(C_i) = x_i$, with $0 \leq x_i \leq 1$, and

$$\sum_{i=1}^n m(C_i) = \sum_{i=1}^n x_i = 1$$

We need to find all variables x_i in accordance with the set of preferences P.

Thus, we get an $m \times n$ linear homogeneous system of equations whose associated matrix is

$$A = (a_{ij}), 1 \leq i \leq m \quad \text{and} \quad 1 \leq j \leq n$$

In order for this system to have non-null solutions, the rank of the matrix A should be strictly less than n.

2.2. Classification of Linear Decision-Making Problems.

a). We say that a linear decision-making problem is consistent if, by any substitution of a variable x_i from an equation into another equation, we get a result in agreement with all equations.

b.) We say that a linear decision-making problem is weakly inconsistent if by at least one substitution of a variable x_i from an equation into another equation we get a result in disagreement with at least another equation in the following ways:

$$\left\{ \begin{array}{l} x_i = k_1 \cdot x_j, \quad k > 1 \\ x_i = k_2 \cdot x_j \quad k_2 > 1, \quad k_2 \neq k_1 \end{array} \right\} \quad (WD1)$$

or

$$\left\{ \begin{array}{l} x_i = k_1 \cdot x_j, \quad 0 < k < 1 \\ x_i = k_2 \cdot x_j \quad 0 < k_2 < 1, \quad k_2 \neq k_1 \end{array} \right\} \quad (WD2)$$

or

$$\{x_i = k \cdot x_j, k \neq 1\} \quad (WD3)$$

(WD1)-(WD3) are weak disagreements, in the sense that for example a variable x is always y , but with different ratios (for example: $x=3y$ and $x=5y$).

All disagreements in this case should be like (WD1)-(WD3).

c). We say that a linear decision-making problem is strongly inconsistent if, by at least one substitution of a variable x_i from an equation into another equation, we get a result in disagreement with at least another equation in the following way:

$$\left\{ \begin{array}{l} x_i = k_1 \cdot x_j; \\ x_i = k_2 \cdot x_j; \end{array} \right\}, \quad (SD4)$$

with $0 < k_1 < 1 < k_2$ or $0 < k_2 < 1 < k_1$ (i.e. from one equation one gets $x_i < x_j$ while from the other equation one gets the opposite inequality: $x_j < x_i$)

At least one inconsistency like (SD4) should exist, no matter if other types of inconsistencies like (WD1)-(WD3) may occur or not.

Compute the determinant of A.

a). If $\det(A)=0$, the decision problem is consistent, since the system of equations is dependent.

It is not necessarily to parameterize the system.

In the case we have parameterized, we can use the Fairness Principle – i.e. setting all parameters equal $\alpha_1 = \alpha_2 = \dots \alpha_n = \alpha_1 > 0$.

Solve this system; find its general solution.

Replace the parameters and secondary variables, getting a particular solution.

Normalize this particular solution (dividing each component by the sum of all components).

We get the priority vector (whose sum of its components should be 1).

b). If $\det A \neq 0$, the decision problem is inconsistent, since the homogeneous linear system has only the null-solution.

b1). If the inconsistency is weak, then parameterize the right-hand side coefficients, and denote the system matrix $A(\alpha)$.

Compute $\det(A(\alpha)) = 0$ in order to get the parametric equation. If the Fairness Principle is used, set all parameters equal, and solve for $\alpha > 0$.

Replace α in $A(\alpha)$ and solve the resulting dependent homogeneous linear system. Similarly as in a), replace each secondary variable by 1, and normalize the particular solution in order to get the priority vector.

b2). If the inconsistency is strong, the Fairness Principle may not work

properly. Another approachable principle might be designed. Or, get more information and revise the strong inconsistencies of the decision-making problem.

2.3. Comparison between AHP and α -D MCDM:

- a). α -D MCDM's general solution includes all particular solutions, that of AHP as well;
- b). α -D MCDM uses all kind of comparisons between criteria, not only pairwise comparisons;
- c). for consistent problems, AHP and α -D MCDM/Fairness-Principle give the same result;
- d). for large inputs, in α -D MCDM we can put the equations under the form of a matrix (depending on some parameters alphas), and then compute the determinant of the matrix which should be zero; after that, solve the system (all can be done on computer using math software); the software such as MATHEMATICA and MAPPLE for example can do these determinants and calculate the solutions of this linear system;
- e). α -D MCDM can work for larger classes of preferences, i.e. preferences that can be transformed in homogeneous linear equations, or in non-linear equations and/or inequalities – see more below.

2.4. Generalization of α -D MCDM

Let each preference be expressed as a linear or non-linear equation or inequality. All preferences together will form a system of linear/non-linear equations/inequalities, or a mixed system of equations and inequalities. Solve this system, looking for a strictly positive solution (i.e. all unknowns $x_i > 0$). Then normalize the solution vector.

If there are more such numerical solutions, do a discussion: analyze the normalized solution vector in each case.

If there is a general solution, extract the best particular solution by replacing the secondary variables by some numbers such that the resulting particular solution is positive, and then normalizing. If there is no strictly positive

solution, parameterize the coefficients of the system, find the parametric equation, and look for some principle to apply in order to find the numerical values of the parameters α 's. A discussion might also be involved. We may get undetermined solutions.

3 Degrees of Consistency and Inconsistency in α -D MCDM/Fairness-Principle

For α -D MCDM/Fairness-Principle in consistent and weak consistent decision-making problems, we have the followings:

- a).** If $0 < \alpha < 1$, then α is the degree of consistency of the decision-making problem, and $\beta = 1 - \alpha$ is the degree of inconsistency of the decision-making problem.
- b).** If $\alpha > 1$, then $1/\alpha$ is the degree of consistency of the decision-making problem, and $\beta = 1 - 1/\alpha$ is the degree of inconsistency of the decision-making problem.

4 Principles of α -D MCDM (Second Part)

- a).** In applications, for the second part of α -D Method, the Fairness Principle can be replaced by other principles.

Expert's Opinion. For example, if we have information that a preference's coefficient should be discounted twice more than another coefficient (due to an expert's opinion), and another preference's coefficient should be discounted a third of another one, then appropriately we set for example: $\alpha_1 = 2\alpha_2$ and respectively $\alpha_3 = [1/3]\alpha_4$, etc. in the parametric equation.

- b).** For α -D/Fairness-Principle or Expert's Opinion. Another idea herein is to set a threshold of consistency t_c (or implicitly a threshold of inconsistency t_i). Then, if the degree of consistency is smaller than a required t_c , the Fairness Principle or Expert's Opinion (whichever was used) should be discharged, and another principle of finding all parameters α 's should be designed; and similarly if the degree of inconsistency is bigger than t_i .

- c).** One may measure the system's accuracy (or error) for the case when all m preferences can be transformed into equations; for example, preference P_i is transformed into an equation $f_i(x_1, x_2, \dots, x_n) = 0$; then we need to find the unknowns $x_1, x_2, \dots, x_n$ such that:

$$e(x_1, x_2, \dots, x_n) = \sum_{i=1}^m |f_i(x_1, x_2, \dots, x_n)|$$

is minimum, where “e(...)” means error.

Calculus theory (partial derivatives) can be used to find the minimum (if this does exist) of a function of n variables, $e(x_1, x_2, \dots, x_n)$, with $e : R_+^n \rightarrow R_+$.

For consistent decision-making problems the system's accuracy/error is zero, so we get the exact result.

We prove this through the fact that the normalized priority vector $[a_1, a_2, \dots, a_n]$, where $x_i = a_i > 0$ for all i , is a particular solution of the system $f_i(x_1, x_2, \dots, x_n) = 0$ for $i = 1, 2, \dots, m$; therefore:

$$\sum_{i=1}^m |f_i(a_1, a_2, \dots, a_n)| = \sum_{i=1}^m |0| = 0$$

But, for inconsistent decision-making problems we find approximations for the variables.

5 Extension of α -D MCDM (Non-Linear α -D MCDM)

It is not difficult to generalize the α -D MCDM for the case when the preferences are non-linear homogeneous (or even non-homogeneous) equations.

This non-linear system of preferences has to be dependent (meaning that its general solution – its main variables - should depend upon at least one secondary variable).

If the system is not dependent, we can parameterize it in the same way. Then, again, in the second part of this Non-Linear α -D MCDM we assign some values to each of the secondary variables (depending on extra-information we might receive), and we also need to design a principle which will help us to find the numerical values for all parameters. We get a particular solution (such extracted from the general solution), which normalized will produce our priority vector.

Yet, the Non-Linear α -D MCDM is more complicated, and depends on each non-linear decision-making problem. Let us see some examples.

6 Consistent Example 1

6.1. We use the α -D MCDM. Let the Set of Preferences be: $\{C1, C2, C3\}$, and The Set of Criteria be:

1. C1 is 4 times as important as C2.
2. C2 is 3 times as important as C3.
3. C3 is one twelfth as important as C1.

Let $m(C1) = x$, $m(C2) = y$, $m(C3) = z$.

The linear homogeneous system associated to this decision-making problem is:

$$\begin{cases} x = 4y \\ y = 3z \\ z = \frac{x}{12} \end{cases}$$

whose associated matrix A_1 is:

$$\begin{pmatrix} 1 & -4 & 0 \\ 0 & 1 & -3 \\ -1/12 & 0 & 1 \end{pmatrix},$$

whence $\det(A_1) = 0$, so the DM problem is consistent.

Solving this homogeneous linear system we get its general solution that we set as a solution vector $[12z \ 3z \ z]$, where z can be any real number (z is considered a secondary variable, while $x=12z$ and $y=3z$ are main variables).

Replacing $z=1$ into the solution vector, the solution vector becomes $[12 \ 3 \ 1]$, and then normalizing (dividing by $12+3+1=16$ each vector component) we get the priority vector: $[12/16 \ 3/16 \ 1/16]$, so the preference will be on C1.

6.2. Using AHP, we get the same result.

The preference matrix is:

$$\begin{pmatrix} 1 & 4 & 12 \\ 1/4 & 1 & 3 \\ 1/12 & 1/3 & 1 \end{pmatrix}$$

whose maximum eigenvalue is $\lambda_{max} = 3$ and its corresponding normalized eigenvector (Perron- Frobenius vector) is $[12/16 \ 3/16 \ 1/16]$.

6.3. Using Mathematica 7.0 Software:

Using MATHEMATICA 7.0 software, we graph the function:


$$h(x, y) = |x - 4y| + |3x + 4y - 3| + |13x + 12y - 12|$$

with $x, y \in [0, 1]$, which represents the consistent decision-making problem's associated system:

$$x/y = 4, y/z = 3, x/z = 12, \quad \text{and} \quad x + y + z = 1, x > 0, y > 0, z > 0$$

$$In[1] := Plot3D[Abs[x - 4y] + Abs[3x + 4y - 3] + Abs[13x + 12y - 12],$$

$$\{x, 0, 1\}, \{y, 0, 1\}]$$



The minimum of this function is zero, and occurs for $x=12/16$, $y=3/16$.

If we consider the original function of three variables associated with $h(x,y)$ we have:

$$H(x, y, z) = |x - 4y| + |y - 3z| + |x - 12z|, x + y + z = 1$$

with $x, y, z \in [0, 1]$, we also get the minimum of $H(x,y,z)$ being zero, which occurs for $x=12/16$, $y=3/16$, $z=1/16$.

7 Weak Inconsistent Examples where AHP does not Work

The Set of Preferences is $\{C1, C2, C3\}$.

7.1. Weak Inconsistent Example 2.

7.1.1 α -D MCDM method.

The Set of Criteria is:

1. C1 is as important as 2 times C2 plus 3 times C3.
2. C2 is half as important as C1.
3. C3 is one third as important as C1.

Let $m(C1) = x$, $m(C2) = y$, $m(C3) = z$;

$$\begin{cases} x = 2y + 3z \\ y = \frac{x}{2} \\ z = \frac{x}{3} \end{cases}$$

AHP cannot be applied on this example because of the form of the first preference, which is not a pairwise comparison.

If we solve this homogeneous linear system of equations as it is, we get $x=y=z=0$, since its associated matrix is:

$$\begin{pmatrix} 1 & -2 & -3 \\ -1/2 & 1 & 0 \\ -1/3 & 0 & 1 \end{pmatrix} = -1 \neq 0$$

but the null solution is not acceptable since the sum $x+y+z$ has to be 1. Let us parameterise each right-hand side coefficient and get the general solution of the above system:

$$\begin{cases} x = 2\alpha_1 y + 3\alpha_2 z & (1) \\ y = \frac{\alpha_3}{2}x & (2) \\ z = \frac{\alpha_4}{3}x & (3) \end{cases}$$

where $\alpha_1, \alpha_2, \alpha_3, \alpha_4 > 0$.

Replacing (2) and (3) in (1) we get

$$x = 2\alpha_1 \left(\frac{\alpha_3}{2}x \right) + 3\alpha_2 \left(\frac{\alpha_4}{3}x \right)$$

$$1 \cdot x = (\alpha_1\alpha_3 + \alpha_1\alpha_4) \cdot x$$

whence

$$\alpha_1\alpha_3 + \alpha_1\alpha_4 = 1 \quad (\text{parametric equation}) \quad (4)$$

The general solution of the system is:

$$\begin{cases} y = \frac{\alpha_3}{2}x \\ z = \frac{\alpha_4}{3}x \end{cases}$$

whence the priority vector:

$$\begin{bmatrix} x & \frac{\alpha_3}{2}x & \frac{\alpha_4}{3}x \end{bmatrix} \rightarrow \begin{bmatrix} 1 & \frac{\alpha_3}{2} & \frac{\alpha_4}{3} \end{bmatrix}$$

Fairness Principle: discount all coefficients with the same percentage: so, replace

$\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = \alpha > 0$ in (4) we get $\alpha^2 + \alpha^2 = 1$, whence $\alpha = \frac{\sqrt{2}}{2}$

Priority vector becomes:

$$\begin{bmatrix} 1 & \frac{\sqrt{2}}{4} & \frac{\sqrt{2}}{6} \end{bmatrix}$$

and normalizing it:

$$\begin{array}{ccc} [0.62923 & 0.22246 & 0.14831] \\ C1 & C2 & C3 \\ x & y & z \end{array}$$

Preference will be on C1, the largest vector component. Let us verify it:

$\frac{y}{x} \cong 0.35354$ instead of 0.50, i.e. $\frac{\sqrt{2}}{2} = 70.71\%$ of the original.

$\frac{z}{x} \cong 0.23570$ instead of 0.33333, i.e. 70.71% of the original.

$x \cong 1.41421y + 2.12132z$ instead of $2y + 3z$, i.e. 70.71% of 2 respectively 70.71% of 3.

So, it was a fair discount for each coefficient.

7.1.2. Using Mathematica 7.0 Software:

Using MATHEMATICA 7.0 software, we graph the function:

$$g(x, y) = |4x - y - 3| + |x - 2y| + |4x + 3y - 3|,$$

with $x, y \in [0, 1]$, which represents the weak inconsistent decision-making problem's associated system:

$$x - 2y - 3z = 0, x - 2y = 0, x - 3z = 0,$$

and


$$x + y + z = 1, x > 0, y > 0, z > 0.$$

by solving $z=1-x-y$ and replacing it in

$$G(x, y, z) = |x - 2y - 3z| + |x - 2y| + |x - 3z|$$

with $x \in [0, 1], y \in [0, 1], z \in [0, 1]$,

`In[2] := Plot3D[Abs[4x - y - 3] + Abs[x - 2y] + Abs[4x + 3y - 3], {x, 0, 1}, {y, 0, 1}]`



Then find the minimum of $g(x,y)$ if any:

In[3]:= FindMinValue

$[Abs[4x - y - 3] + Abs[x - 2y] + Abs[4x + 3y - 3], x + y \leq 1, x > 0, y > 0, x, y]$

The following result is returned:

Out[3]:= 0.841235.

FindMinValue::eit: The algorithm does not converge to the tolerance of $4.806217383937354 \cdot 10^{-6}$ in 500 iterations. The best estimated solution, with feasibility residual, KKT residual, or complementary residual of $\{0.0799888, 0.137702, 0.0270028\}$, is returned.

7.1.3. Matrix Method of using α -Discounting.

The determinant of the homogeneous linear system (1), (2), (3) is:

$$\begin{vmatrix} 1 & -2\alpha_1 & -3\alpha_2 \\ -\frac{1}{2}\alpha_3 & 1 & 0 \\ -\frac{1}{2}\alpha_4 & 0 & 1 \end{vmatrix} = (1 + 0 + 0) - (\alpha_2\alpha_4 + \alpha_1\alpha_3) = 0$$

or $\alpha_1\alpha_3 + \alpha_2\alpha_4 = 1$ (parametric equation).

The determinant has to be zero in order for the system to have non-null solutions.

The rank of the matrix is 2.

So, we find two variables, for example it is easier to solve for y and z from the last two equations, in terms of x :

$$\begin{cases} y = \frac{1}{2}\alpha_3x \\ z = \frac{1}{3}\alpha_4x \end{cases}$$

and the procedure follows the same steps as in the previous one.

Let us change Example 1 in order to study various situations.

7.2. Weak Inconsistent Example 3, which is more weakly inconsistent than Example 2.

1. Same as in Example 2.
2. C_2 is 4 times as important as C_1

3. Same as in Example 2.

$$\begin{cases} x = 2\alpha_1 y + 3\alpha_2 z \\ y = 4\alpha_3 x \\ z = \frac{\alpha_4}{3} x \end{cases}$$

$$x = 2\alpha_1 (4\alpha_3 x) + 3\alpha_2 \left(\frac{\alpha_4}{3} \right) x$$

$$1 \cdot x = (8\alpha_1\alpha_3 + \alpha_2\alpha_4) \cdot x$$

$$8\alpha_1\alpha_3 + \alpha_2\alpha_4 = 1(\text{parametric equation})$$

$$\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = \alpha > 0$$

$$9\alpha^2 = 1 \Rightarrow \alpha = \frac{1}{3}$$

$$\begin{bmatrix} x & 4\alpha_3 x & \frac{\alpha_4}{3} x \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 4\alpha_3 & \frac{\alpha_4}{3} \end{bmatrix}$$

$$\begin{bmatrix} 1 & \frac{4}{3} & \frac{1}{9} \end{bmatrix} = \begin{bmatrix} \frac{9}{9} & \frac{12}{9} & \frac{1}{9} \end{bmatrix}$$

normalized:

$$\begin{bmatrix} \frac{9}{22} & \frac{12}{22} & \frac{1}{22} \end{bmatrix}$$

$\frac{y}{x} = 1.333$ instead of 4;

$\frac{z}{x} = 0.111$ instead of 0.333;

$x = 0.667y + 1 \cdot z$ instead of $2y + 3z$.

Each coefficient was reduced at $\frac{1}{3}$ ($= 33.33\%$)

The bigger is the inconsistency ($\beta \rightarrow 1$), the bigger is the discounting ($\alpha \rightarrow 0$).

7.3. Weak Inconsistent Example 4, which is even more inconsistent than Example 3.

1. Same as in Example 2.

2. Same as in Example 3.

3. C 3 is 5 times as important as C1

$$\begin{cases} x = 2\alpha_1 y + 3\alpha_2 z \\ y = 4\alpha_3 x \\ z = 5\alpha_4 x \end{cases}$$

$$x = 2\alpha_1 (4\alpha_3 x) + 3\alpha_2 (5\alpha_4 x)$$

$$1 \cdot = (8\alpha_1\alpha_3 + 15\alpha_2\alpha_4) x$$

whence $8\alpha_1\alpha_3 + 15\alpha_2\alpha_4 = 1$ (parametric equation)

$$\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = \alpha > 0, 23\alpha^2 = 1, \alpha = \frac{\sqrt{23}}{23}$$

Normalized: [0.34763 0.28994 0.36243]

$\frac{y}{x} \cong 0.83405$ instead of 4, i.e. reduced at $\frac{\sqrt{23}}{23} = 20.85\%$;

$\frac{z}{x} \cong 1.04257$ instead of 5;

$x \cong 0.41703y + 0.62554 \cdot z$ instead of $2x + 3y$.

Each coefficient was reduced at $\alpha = \frac{\sqrt{23}}{23} \cong 20.85\%$

7.4. Consistent Example 5.

When we get $\alpha = 1$, we have a consistent problem.

Suppose the preferences:

1. Same as in Example 2.
2. C 2 is one fourth as important as C1 .
3. C 3 is one sixth as important as C1 .

The system is:

$$\begin{cases} x = 2y + 3z \\ y = \frac{x}{4} \\ z = \frac{x}{6} \end{cases}$$

7.4.1. First Method of Solving this System.

Replacing the second and third equations of this system into the first, we get:

$$x = 2 \left(\frac{x}{4} \right) + 3 \left(\frac{x}{6} \right) = \frac{x}{2} + \frac{x}{2} = x$$

which is an identity (so, no contradiction).

General solution:

$$\begin{bmatrix} x & \frac{x}{4} & \frac{x}{6} \end{bmatrix}$$

Priority vector

$$\begin{bmatrix} 1 & \frac{1}{4} & \frac{1}{6} \end{bmatrix}$$

Normalized is

$$\begin{bmatrix} \frac{12}{17} & \frac{3}{17} & \frac{2}{17} \end{bmatrix}$$

7.4.2. Second Method of Solving this System.

Let us parameterize:

$$\begin{cases} x = 2\alpha_1 y + 3\alpha_2 z \\ y = \frac{\alpha_3}{4} x \\ z = \frac{\alpha_4}{6} x \end{cases}$$

Replacing the last two equations into the first we get:

$$x = 2\alpha_1 \left(\frac{\alpha_3}{4} x \right) + 3\alpha_2 \left(\frac{\alpha_4}{6} x \right) = \frac{\alpha_1 \alpha_3}{2} x + \frac{\alpha_2 \alpha_4}{2} x$$

$$1 \cdot x = \frac{\alpha_1 \alpha_3 + \alpha_2 \alpha_4}{2} \cdot x$$

whence $1 = \frac{\alpha_1 \alpha_3 + \alpha_2 \alpha_4}{2}$ or $\alpha_1 \alpha_3 + \alpha_2 \alpha_4 = 2$.

Consider the fairness principle: $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = \alpha > 0$, then $2\alpha^2 = 2$, $\alpha = \pm 1$, but we take only the positive value $\alpha = 1$ (as expected for a consistent problem).

Let us check:

$$\frac{y}{x} = \frac{\frac{3}{17}}{\frac{12}{17}} = \frac{1}{4}, \text{ exactly as in the original system;}$$

$$\frac{z}{x} = \frac{\frac{2}{17}}{\frac{12}{17}} = \frac{1}{6}, \text{ exactly as in the original system;}$$

$$x = 2y + 3z \text{ since } z = 2 \left(\frac{x}{4} \right) + 3 \left(\frac{x}{6} \right);$$

hence all coefficients were left at $\alpha = 1$ (=100%) of the original ones.

No discount was needed.

7.5. General Example 6.

Let us consider the general case:

$$\begin{cases} x = a_1y + a_2z \\ y = a_3x \\ z = a_4x \end{cases}$$

where $a_1, a_2, a_3, a_4 > 0$.

Let us parameterize:

$$\begin{cases} x = a_1\alpha_1y + a_2\alpha_2z \\ y = a_3\alpha_3x \\ z = a_4\alpha_4x \end{cases}$$

with $\alpha_1, \alpha_2, \alpha_3, \alpha_4 > 0$.

Replacing the second and third equations into the first, we get:

$$x = a_1\alpha_1(a_3\alpha_3x) + a_2\alpha_2(a_4\alpha_4x)$$

$$x = a_1a_3\alpha_1\alpha_3x + a_2a_4\alpha_2\alpha_4x$$

whence $a_1a_3\alpha_1\alpha_3x + a_2a_4\alpha_2\alpha_4 = 1$ (parametric equation)

The general solution of the system is: $(x, a_3\alpha_3x, a_4\alpha_4x)$

The priority vector is $\begin{bmatrix} 1 & a_3\alpha_3 & a_4\alpha_4 \end{bmatrix}$

Consider the fairness principle: $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = \alpha > 0$, we get:

$$\alpha^2 = \frac{1}{a_1a_3 + a_2a_4},$$

so

$$\alpha = \frac{1}{\sqrt{a_1a_3 + a_2a_4}}$$

i). if $\alpha \in [0, 1]$, then α is the degree of consistency of the problem, while $\beta = 1 - \alpha$ is the degree of the inconsistency of the problem.

ii). If $\alpha > 1$, then $\frac{1}{\alpha}$ is the degree of consistency, while $\beta = 1 - \frac{1}{\alpha}$ is the degree of α inconsistency.

When the degree of consistency $\rightarrow 0$, the degree of inconsistency $\rightarrow 1$, and reciprocally.

Discussion of the General Example 6.

Suppose the coefficients a_1, a_2, a_3, a_4 become big such that $a_1a_3 + a_2a_4 \rightarrow \infty$,

then $\alpha \rightarrow 0$, and $\beta \rightarrow 1$.

Particular Example 7.

Let us see a particular case when a_1, a_2, a_3, a_4 make $a_1a_3 + a_2a_4$ big:

$$a_1 = 50, a_2 = 20, a_3 = 100, a_4 = 250,$$

then $\alpha = \frac{1}{\sqrt{50 \cdot 100 + 20 \cdot 250}} = \frac{1}{\sqrt{10000}} = \frac{1}{100} = 0.01 =$ degree of consistency,
whence $\beta = 0.99$ degree of inconsistency.

The priority vector for Particular Example 7 is $[1 \ 100(0.01) \ 250(0.01)] = [1 \ 1 \ 2.5]$ which normalized is:

$$\left[\frac{2}{9} \quad \frac{2}{9} \quad \frac{5}{9} \right]$$

Particular Example 8.

Another case when a_1, a_2, a_3, a_4 make the expression $a_1a_3 + a_2a_4$ a tiny positive number: $a_1 = 0.02, a_2 = 0.05, a_3 = 0.03, a_4 = 0.02$, then

$$\alpha = \frac{1}{\sqrt{0.02 \cdot (0.03) + 0.05 \cdot (0.02)}} = \frac{1}{0.04} = 25 > 1$$

Then $\frac{1}{\alpha} = \frac{1}{25} = 0.04$ is the degree of consistency of the problem, and 0.96 the degree of inconsistency.

The priority vector for Particular Example 8 is

$$\left[1 \quad a_3\alpha \quad a_4\alpha \right] = \left[1 \quad 0.03(25) \quad 0.02(25) \right] = \left[1 \quad 0.75 \quad 0.50 \right]$$

which normalized is

$$\left[\frac{4}{9} \quad \frac{3}{9} \quad \frac{3}{9} \right]$$

Let us verify:

$\frac{y}{x} = \frac{3}{9} \div \frac{4}{9} = 0.75$ instead of 0.03, i.e. $\alpha = 25$ times larger (or 2500%);
 $\frac{z}{x} = \frac{2}{9} \div \frac{4}{9} = 0.50$ instead of $x = 0.02y + 0.05z$ (0.50 is 25 times larger than 0.02 and 1.25 is 25 times larger than 0.05) because $\frac{4}{9} = 0.50 \left(\frac{3}{9}\right) + 1.25 \left(\frac{2}{9}\right)$

8.1. Jean Dezert's Weak Inconsistent Example 9.

Let $\alpha_1, \alpha_2, \alpha_3 > 0$ be the parameters. Then:

$$\left\{ \begin{array}{l} \frac{x}{y} = 3\alpha_1 \\ \frac{x}{z} = 4\alpha_2 \\ \frac{y}{z} = 5\alpha_3 \end{array} \right. \quad \begin{array}{l} (5) \\ (6) \\ (7) \end{array} \Rightarrow \frac{x}{y} \cdot \frac{x}{z} = (3\alpha_1) \cdot (4\alpha_2) \Rightarrow \frac{y}{z} = 12\alpha_1\alpha_2$$

On order for $\frac{y}{z} = 12\alpha_1\alpha_2$ we need to have $12\alpha_1\alpha_2 = 5\alpha_3$ or

$$2.4\alpha_1\alpha_2 = \alpha_3 (\text{ParametricEquation}) \quad (8)$$

Solving this system:

$$\left\{ \begin{array}{l} \frac{x}{y} = 3\alpha_1 \Rightarrow y = 3\alpha_1 \cdot x \\ \frac{x}{z} = 4\alpha_2 \Rightarrow x = 4\alpha_2 \cdot z \\ \frac{y}{z} = 5\alpha_3 \Rightarrow y = 12\alpha_1\alpha_2 z \end{array} \right.$$

we get the general solution:

$$\left[\begin{array}{ccc} 4\alpha_2 z & 5(2.4\alpha_1\alpha_2) & z \end{array} \right]$$

$$\left[\begin{array}{ccc} 4\alpha_2 z & 12\alpha_1\alpha_2 z & z \end{array} \right]$$

General normalized priority vector is:

$$\left[\begin{array}{ccc} \frac{4\alpha_2}{4\alpha_2+12\alpha_1\alpha_2+1} & \frac{12\alpha_1\alpha_2}{4\alpha_2+12\alpha_1\alpha_2+1} & \frac{1}{4\alpha_2+12\alpha_1\alpha_2+1} \end{array} \right]$$

where $\alpha_1\alpha_2 > 0$; ($\alpha_3 = 2.4\alpha_1\alpha_2$)

Which α_1 and α_2 give the best result? How to measure it? This is the greatest challenge!

α -Discounting Method includes all solutions (all possible priority vectors which make the matrix consistent).

Because we have to be consistent with all proportions (i.e. using the Fairness Principle of finding the parameters' numerical values), there should be the same discounting of all three proportions (5), (6), and (7), whence

$$\alpha_1 = \alpha_2 = \alpha_3 > 0 \quad (9)$$

The parametric equation (8) becomes $2.4\alpha_1^2 = \alpha_1$ or $2.4\alpha_1^2 - \alpha_1 = 0$ or $\alpha_1(2.4\alpha_1 - 1) = 0$ whence $\alpha_1 = 0$ or $\alpha_1 = \frac{1}{2.4} = \frac{5}{12}$
 $\alpha_1 = 0$ is not good, contradicting (9). Our system becomes now:

$$\left\{ \begin{array}{l} \frac{y}{x} = 3 \cdot \frac{5}{12} = \frac{15}{12} \\ \frac{x}{z} = 4 \cdot \frac{5}{12} = \frac{20}{12} \\ \frac{y}{z} = 5 \cdot \frac{5}{12} = \frac{25}{12} \end{array} \right. \quad \begin{array}{l} (10) \\ (11) \\ (12) \end{array}$$

We see that (10) and (11) together give

$$\frac{y}{x} \cdot \frac{x}{z} = \frac{15}{12} \text{ or } \frac{y}{z} = \frac{25}{12}$$

so, they are now consistent with (12).

From (11) we get $x = \frac{20}{12}z$ and from (12) we get $y = \frac{25}{12}z$

The priority vector is:

$$\left[\frac{20}{12}z \quad \frac{25}{12}z \quad 1z \right]$$

which is normalized to:

$$\begin{aligned} \frac{\frac{20}{12}}{\frac{20}{12} + \frac{25}{12} + 1} &= \frac{\frac{20}{12}}{\frac{20}{12} + \frac{25}{12} + \frac{12}{12}} = \frac{20}{57}, \frac{\frac{25}{12}}{\frac{20}{12} + \frac{25}{12} + \frac{12}{12}} = \frac{25}{57}, \frac{1}{\frac{20}{12} + \frac{25}{12} + \frac{12}{12}} = \frac{12}{57}, i.e. \\ &\begin{matrix} C_1 & C_2 & C_3 \\ \left[\frac{20}{57} & \frac{25}{57} & \frac{12}{57} \right]^T \end{matrix} \\ &\cong \begin{matrix} C_1 & C_2 & C_3 \\ \left[0.3509 & 0.4386 & 0.2105 \right]^T \end{matrix} \\ &\quad \quad \quad \begin{matrix} \uparrow \\ \text{the highest priority} \end{matrix} \end{aligned} \quad (13)$$

Let us study the result:

$$\begin{matrix} C_1 & C_2 & C_3 \\ \left[\frac{20}{57} & \frac{25}{57} & \frac{12}{57} \right]^T \\ x & y & z \end{matrix}$$

<i>Ratios :</i>	Percentage of Discounting :
$\frac{y}{x} = \frac{\frac{25}{57}}{\frac{20}{57}} = \frac{25}{20} = 1.25$ instead of 3;	$\frac{\frac{25}{20}}{3} = \frac{5}{12} = \alpha_1 = 41.\overline{6}\%$
$\frac{x}{z} = \frac{\frac{20}{57}}{\frac{12}{57}} = \frac{20}{12} = \frac{5}{3} = 1.\overline{6}$ instead of 4;	$\frac{\frac{20}{12}}{4} = \frac{5}{12} = \alpha_1 = 41.\overline{6}\%$
$\frac{y}{z} = \frac{\frac{25}{57}}{\frac{12}{57}} = \frac{25}{12} = 2.\overline{083}$ instead of 5;	$\frac{\frac{25}{12}}{5} = \frac{5}{12} = \alpha_1 = 41.\overline{6}\%$

Hence all original proportions, which were respectively equal to 3, 4, and 5 in the 5 problem, were reduced by multiplication with the same factor $\alpha_1 = \frac{5}{12}$, i.e. by getting 41.6% of each of them.

So, it was fair to reduce each factor to the same percentage 41. $\overline{6}\%$ of itself. But this is not the case in Saaty's method: its normalized priority vector is

$$\begin{bmatrix} C_1 & C_2 C_3 \\ 0.2797 & 0.6267 & 0.0936 \end{bmatrix}^T$$

$$\begin{matrix} x & y & z \end{matrix}$$

where

<i>Ratios :</i>	Percentage of Discounting :
$\frac{y}{x} = \frac{0.6267}{0.2797} \cong 2.2406$ instead of 3;	$\frac{2.2406}{3} \cong 74.6867\%$
$\frac{x}{z} = \frac{0.2797}{0.0936} \cong 2.9882$ instead of 4;	$\frac{2.9882}{4} \cong 74.7050\%$
$\frac{y}{z} = \frac{0.6267}{0.0936} \cong 6.6955$ instead of 5;	$\frac{6.6955}{5} \cong 133.9100\%$

Why, for example, the first proportion, which was equal to 3, was discounted to 74.6867% of it, while the second proportion, which was equal to 4, was discounted to another percentage (although close) 74.7050% of it?

Even more doubt we have for the third proportion's coefficient, which was equal to 5, but was increased to 133.9100% of it, while the previous two proportions were decreased; what is the justification for these?

That is why we think our α -D/Fairness-Principle is better justified.

We can solve this same problem using matrices. (5), (6), (7) can be written in another way to form a linear parameterized homogeneous system:

$$\begin{cases} 3\alpha_1 x - y = 0 \\ x - 4\alpha_2 z = 0 \\ y - 5\alpha_3 z = 0 \end{cases} \quad (14)$$

Whose associated matrix is:

$$P_1 = \begin{bmatrix} 3\alpha_1 & -1 & 0 \\ 1 & 0 & -4\alpha_2 \\ 0 & 1 & -5\alpha_3 \end{bmatrix} \quad (15)$$

a). If $\det(P_1) \neq 0$ then the system (10) has only the null solution $x = y = z = 0$.

b) Therefore, we need to have $\det(P_1) = 0$, or $(3\alpha_1)(4\alpha_2) - 5\alpha_3 = 0$, or $2.4\alpha_1\alpha_2 - \alpha_3 = 0$, so we get the same parametric equation as (8).

In this case the homogeneous parameterized linear system (14) has a triple infinity of solutions.

This method is an extension of Saaty's method, since we have the possibility to manipulate the parameters α_1, α_2 , and α_3 . For example, if a second

source tells us that $\frac{x}{z}$ has to be discounted 2 times as much as $\frac{y}{x}$, and $\frac{y}{z}$ should be discounted 3 times less than $\frac{y}{x}$, then we set $\alpha_2 = 2\alpha_1$, and respectively $\alpha_3 = \frac{\alpha_1}{3}$, and the original (5), (6), (7) system becomes:

$$\begin{cases} \frac{y}{x} = 3\alpha_1 \\ \frac{x}{z} = 4\alpha_2 = 4(2\alpha_1) = 8\alpha_1 \\ \frac{y}{z} = 5\alpha_3 = 5\left(\frac{\alpha_1}{3}\right) = \frac{5}{3}\alpha_1 \end{cases} \quad (16)$$

and we solve it in the same way.

8.2. Weak Inconsistent Example 10.

Let us complicate Jean Dezert's Weak Inconsistent Example 9. with one more preference: C_2 is 1.5 times as much as C_1 and C_3 together. The new system is:

$$\begin{cases} \frac{y}{x} = 3 \\ \frac{x}{z} = 4 \\ \frac{y}{z} = 5 \\ y = 1.5(x + z) \\ x, y, z \in [0, 1] \\ x + y + z = 1 \end{cases} \quad (17)$$

We parameterized it:

$$\begin{cases} \frac{y}{x} = 3\alpha_1 \\ \frac{x}{z} = 4\alpha_2 \\ \frac{y}{z} = 5\alpha_3 \\ y = 1.5\alpha_4(x + z) \\ x, y, z \in [0, 1] \\ x + y + z = 1 \end{cases} \quad (18)$$

$$\alpha_1, \alpha_2, \alpha_3, \alpha_4 > 0$$

Its associated matrix is:

$$\begin{bmatrix} 3\alpha_1 & -1 & 0 \\ 1 & 0 & -4\alpha_2 \\ 0 & 1 & -5\alpha_3 \\ 1.5\alpha_4 & -1 & 1.5\alpha_4 \end{bmatrix} \quad (19)$$

The rank of matrix P_2 should be strictly less than 3 in order for the system (18) to have non-null solution.

If we take the first three rows in (19) we get the matrix P_1 , whose determinant should be zero, therefore one also gets the previous parametric equation $2.4\alpha_1\alpha_2 = \alpha_3$.

If we take rows 1, 3, and 4, since they all involve the relations between C_2 and the other criteria C_1 and C_3 we get

$$\begin{bmatrix} 3\alpha_1 & -1 & 0 \\ 0 & 1 & -5\alpha_3 \\ 1.5\alpha_4 & -1 & 1.5\alpha_4 \end{bmatrix} \quad (20)$$

whose determinant should also be zero:

$$\begin{aligned} \det(P_3) &= [3\alpha_1(1.5\alpha_4) + 5\alpha_3(1.5\alpha_4) + 0] - [0 + 3\alpha_1(5\alpha_3) + 0] = \\ &= 4.5\alpha_1\alpha_4 + 7.5\alpha_3\alpha_4 - 15\alpha_1\alpha_3 = 0 \end{aligned} \quad (21)$$

If we take

$$P_4 = \begin{bmatrix} 1 & 0 & -4\alpha_2 \\ 0 & 1 & -5\alpha_3 \\ 1.5\alpha_4 & -1 & 1.5\alpha_4 \end{bmatrix} \quad (22)$$

then

$$\begin{aligned} \det(P_4) &= [1.5\alpha_4 + 0 + 0] - \\ &- [-6\alpha_2\alpha_4 + 5\alpha_3 + 0] = 1.5\alpha_4 + 6\alpha_2\alpha_4 - 5\alpha_3 = 0 \end{aligned} \quad (23)$$

If we take

$$P_5 = \begin{bmatrix} 3\alpha_1 & -1 & 0 \\ 1 & 0 & -4\alpha_2 \\ 1.5\alpha_4 & -1 & 1.5\alpha_4 \end{bmatrix} \quad (24)$$

then

$$\begin{aligned} \det(P_5) &= [0 + 0 + 6\alpha_2\alpha_4] - \\ &- [0 + 12\alpha_1\alpha_2 - 1.5\alpha_4] = +6\alpha_2\alpha_4 - 12\alpha_1\alpha_2 + 1.5\alpha_4 = 0 \end{aligned} \quad (25)$$

So, these four parametric equations form a parametric system:

$$\begin{cases} 2.4\alpha_1\alpha_2 - \alpha_3 = 0 \\ 4.5\alpha_1\alpha_2 + 7.5\alpha_3\alpha_4 - 15\alpha_1\alpha_3 = 0 \\ 1.5\alpha_4 + 6\alpha_2\alpha_4 - 5\alpha_3 = 0 \\ 6\alpha_2\alpha_4 - 12\alpha_1\alpha_2 + 1.5\alpha_4 = 0 \end{cases} \quad (26)$$

which should have a non-null solution.

If we consider $\alpha_1 = \alpha_2 = \alpha_3 = \frac{5}{12} > 0$ as we got at the beginning, then substituting all α 's 12 into the last three equations of the system (26) we get:

$$4.5 \left(\frac{5}{12} \right) \alpha_4 + 7.5 \left(\frac{5}{12} \right) \alpha_4 - 15 \left(\frac{5}{12} \right) \left(\frac{5}{12} \right) = 0 \Rightarrow \alpha_4 = 0.5208\bar{3} = \frac{25}{48}$$

$$1.5\alpha_4 + 6 \left(\frac{5}{12} \right) \alpha_4 - 5 \left(\frac{5}{12} \right) = 0 \Rightarrow \alpha_4 = 0.5208\bar{3}$$

$$6 \left(\frac{5}{12} \right) \alpha_4 - 12 \left(\frac{5}{12} \right) \left(\frac{5}{12} \right) + 1.5\alpha_4 = 0 \Rightarrow \alpha_4 = 0.5208\bar{3}$$

α_4 could not be equal to $\alpha_1 = \alpha_2 = \alpha_3$ since it is an extra preference, because the number of rows was bigger than the number of columns.

So the system is consistent, having the same solution as previously, without having added the fourth preference $y = 1.5 (x + z)$.

9.1. Jean Dezert's Strong Inconsistent Example 11.

9.1.1. The preference matrix is:

$$M_1 = \begin{pmatrix} 1 & 9 & \frac{1}{9} \\ \frac{1}{9} & 1 & 9 \\ 9 & \frac{1}{9} & 1 \end{pmatrix}$$

so,

$$\begin{cases} x = 9y, & x > y \\ x = \frac{1}{9}z, & x < z \\ y = 9z, & y > z \end{cases}$$

The other three equations: $y = \frac{1}{9}x, z = 9x, z = \frac{1}{9}y$ result directly from the previous three ones, so we can eliminate them. From $x \succ y$ and $y \succ z$ (first and third above inequalities) we get $x \succ z$, but the second inequality tells us the opposite: $x \prec z$; that is why we have a strong contradiction/inconsistency. Or, if we combine all three we have $x \succ y \succ z \succ x \dots$ strong contradiction again.

Parameterize:

$$\begin{cases} x = 9\alpha_1 y & (27) \\ x = \frac{1}{9}\alpha_2 z & (28) \\ y = 9\alpha_3 z & (29) \end{cases}$$

where $\alpha_1, \alpha_2, \alpha_3 > 0$.

From (27) we get: $y = \frac{1}{9\alpha_1}x$, from (28) we get $z = \frac{1}{9\alpha_2}x$, which is replaced in (29) and we get: $y = 9\alpha_3 \left(\frac{1}{9\alpha_2}x \right) = \frac{81\alpha_3}{\alpha_2}x$.

So $\frac{1}{9\alpha_1}x = \frac{81\alpha_3}{\alpha_2}x$ or $\alpha_2 = 729\alpha_1\alpha_3$ (parametric equation).

The general solution of the system is:

$$\left(x, \frac{1}{9\alpha_1}x, \frac{9}{\alpha_2}x \right)$$

The general priority vector is:

$$\left[1 \quad \frac{1}{9\alpha_1} \quad \frac{9}{\alpha_2} \right]$$

Consider the fairness principle, then $\alpha_1 = \alpha_2 = \alpha_3 = \alpha > 1$ are replaced into the parametric equation: $\alpha = 729\alpha^2$, whence $\alpha = \frac{1}{729} = \frac{1}{9^3}$.

The particular priority vector becomes $\left[1 \quad 9^2 \quad 9^4 \right] = \left[1 \quad 81 \quad 6561 \right]$ and normalized

$$\left[\frac{1}{6643} \quad \frac{81}{6643} \quad \frac{6561}{6643} \right]$$

Because the consistency is $\alpha = \frac{1}{729} = 0.00137$ is extremely low, we can disregard this solution (and the inconsistency is very big $\beta = 1 - \alpha = 0.99863$).

9.1.2. Remarks.

a). If in M_1 we replace all six 9's by a bigger number, the inconsistency of the system will increase. Let us use 11. 1

Then $\alpha = \frac{1}{11^3} = 0.00075$ (consistency), while inconsistency $\beta = 0.99925$.

b). But if in M_1 we replace all 9's by the smaller positive number greater than 1, the 1 consistency decreases. Let us use 2. Then $\alpha = \frac{1}{2^3} = 0.125$ and

$$\beta = 0.875$$

c). Consistency is 1 when replacing all six 9's by 1.

d). Then, replacing all six 9's by a positive sub unitary number, consistency 1 decreases again. For example, replacing by 0.8 we get $\alpha = \frac{1}{0.8^3} = 1.953125 > 1$, whence $\frac{1}{\alpha} = 0.512$ (consistency) and $\beta = 0.488$ (inconsistency).

9.2. Jean Dezert's Strong Inconsistent Example 12.

The preference matrix is:

$$M_2 = \begin{pmatrix} 1 & 5 & \frac{1}{5} \\ \frac{1}{5} & 1 & 5 \\ 5 & \frac{1}{5} & 1 \end{pmatrix}$$

which is similar to M_1 where we replace all six 9's by 5's.

$\alpha = \frac{1}{5} = 0.008$ (consistency) and $\beta = 0.992$ (inconsistency)

The priority vector is $[1 \ 5^2 \ 5^4] = [1 \ 25 \ 625]$ and normalized

$$[\frac{1}{651} \ \frac{25}{651} \ \frac{625}{651}]$$

M_2 is a little more consistent than M_1 because $0.00800 < 0.00137$, but still not enough, so this result is also discarded.

9.3. Generalization of Jean Dezert's Strong Inconsistent Examples.

General Example 13.

Let the preference matrix be:

$$M_t = \begin{pmatrix} 1 & t & \frac{1}{t} \\ \frac{1}{t} & 1 & t \\ t & \frac{1}{t} & 1 \end{pmatrix}$$

with $t > 0$, and $c(M_t)$ the consistency of M_t , $y(M_t)$ inconsistency of M_t .

We have for the Fairness Principle:

$$\lim_{t \rightarrow 1} c(M_t) = 1 \quad \text{and} \quad \lim_{t \rightarrow 1} i(M_t) = 0$$

$$\lim_{t \rightarrow +\infty} c(M_t) = 0 \quad \text{and} \quad \lim_{t \rightarrow +\infty} i(M_t) = 1$$

$$\lim_{t \rightarrow 0} c(M_t) = 0 \quad \text{and} \quad \lim_{t \rightarrow 0} i(M_t) = 1$$

Also $\alpha = \frac{1}{t^3}$, the priority vector is $\begin{bmatrix} 1 & t^2 & t^4 \end{bmatrix}$ which is normalized as

$$\begin{bmatrix} \frac{1}{1+t^2+t^4} & \frac{t^2}{1+t^2+t^4} & \frac{t^4}{1+t^2+t^4} \end{bmatrix}$$

In such situations, when we get strong contradiction of the form $x_i y_j z_i x$ or similarly $x_j z_i x$, etc. and the consistency is tiny, we can consider that $x=y=z=1/3$ (so no criterion is preferable to the other – as in Saaty's AHP), or just $x+y+z=1$ (which means that one has the total ignorance too: $C1 \cup C2 \cup C3$).

10 Strong Inconsistent Example 14

Let $C = \{C1, C2\}$, and $P = \{C1 \text{ is important twice as much as } C2; C2 \text{ is important 5 times as much as } C1\}$. Let $m(C1)=x$, $m(C2)=y$. Then: $x=2y$ and $y=5x$ (it is a strong inconsistency since from the first equation we have $x_i y_j$, while from the second $y_i x_j$).

Parameterize: $x = 2\alpha_1 y$, $y = 5\alpha_2 x$, whence we get $2\alpha_1 = 1/(5\alpha_2)$, or $10\alpha_1\alpha_2 = 1$

If we consider the Fairness Principle, then $\alpha_1 = \alpha_2 = \alpha > 0$, and one gets $\alpha = \frac{\sqrt{10}}{10} \approx 31.62\%$ consistency; priority vector is $[0.39 \ 0.61]$, hence $y_i x_j$. An explanation can be done as in paraconsistent logic (or as in neutrosophic logic): we consider that the preferences were honest, but subjective, therefore it is possible to have two contradictory statements true simultaneously since a criterion $C1$ can be more important from a point of view than $C2$, while from another point of view $C2$ can be more important than $C1$. In our decision- making problem, not having any more information and having rapidly being required to take a decision, we can prefer $C2$, since $C2$ is 5 times more important than $C1$, while $C1$ is only 2 times more important than $C2$, and $5_i 2_j$.

If it's no hurry, more prudent would be in such dilemma to search for more information on $C1$ and $C2$.

If we change Example 14 under the form: $x=2y$ and $y=2x$ (the two coefficients set equal), we get $\alpha = \frac{1}{2}$, so the priority vector is $[0.5 \ 0.5]$ and decision-making problem is undecidable.

11 Non-Linear Equation System Example 15

Let $= \{C1, C2, C3\}$, $m(C1)=x$, $m(C2)=y$, $m(C3)=z$. Let F be:

1. $C1$ is twice as much important as the product of $C2$ and $C3$.
2. $C2$ is five times as much important as $C3$.

We form the non-linear system: $x=2yz$ (non-linear equation) and $y=5z$ (linear equation).

The general solution vector of this mixed system is: $[10z^2 \ 5z \ z]$, where $z \geq 0$. A discussion is necessary now.

a). You see for sure that $y \leq z$, since $5z \leq z$ for z strictly positive. But we don't see anything what the position of x would be?

b). Let us simplify the general solution vector by dividing each vector component by $z \geq 0$, thus we get: $[10z \ 5 \ 1]$.

If $z \in (0, 0.1)$, then $y \leq z \leq x$.

If $z=0.1$, then $y \leq z=x$.

If $z \in (0.1, 0.5)$, then $y \leq x \leq z$.

If $z=0.5$, then $y=x \leq z$.

If $z \geq 0.5$, then $x \leq y \leq z$.

12 Non-Linear/Linear Equation/Inequality Mixed System Example 16

Since in the previous Example 15 has many variants, assume that a new preference comes in (in addition to the previous two preferences): 3. $C1$ is less important than $C3$. The mixed system becomes now: $x=2yz$ (non-linear equation), $y=5z$ (linear equation), and $x \leq z$ (linear inequality). The general solution vector of this mixed system is: $[10z^2 \ 5z \ z]$, where $z \geq 0$ and $10z^2 \leq z$. From the last two inequalities we get $z \in (0, 0.1)$. Whence the priorities are: $y \leq z \leq x$.

13 Future Research

To investigate the connection between α -D MCDM and other methods, such as: the technique for order preference by similarity to ideal solution (TOPSIS) method, the simple additive weighting (SAW) method, Borda-Kendall (BK) method for aggregating ordinal preferences, and the cross-efficiency evaluation method in data envelopment analysis (DEA).

14 Conclusion

We have introduced a new method in the multi-criteria decision making, α - Discounting MCDM. In the first part of this method, each preference is transformed into a linear or non-linear equation or inequality, and all together form a system that is resolved – one finds its general solution, from which one extracts the positive solutions. If the system has only the null solution, or it is inconsistent, then one parameterizes the coefficients of the system.

In the second part of the method, one chooses a principle for finding the numerical values of the parameters {we have proposed herein the Fairness Principle, or Expert's Opinion on Discounting, or setting a Consistency (or Inconsistency) Threshold}. Acknowledgement: The author wants to thank Mr. Atilla Karaman, from the Operations Research Department of KHO SAVBEN in Ankara, Turkey, Ph D student using this α -Discounting Method for Multi-Criteria Decision Making in his Ph D thesis, for his remarks on this paper.

REFERENCES

- [1] J. Barzilai, *Notes on the Analytic Hierarchy Process*, Proc. of the NSF Design and Manufacturing Research Conf., pp. 1–6, Tampa, Florida, January 2001.
- [2] V. Belton, A.E. Gear, *On a Short-coming of Saaty's Method of Analytic Hierarchies*, Omega, Vol. 11, No. 3, pp. 228–230, 1983.
- [3] M. Beynon, B. Curry, P.H. Morgan, *The Dempster-Shafer theory of evidence: An alternative approach to multicriteria decision modeling*, Omega, Vol. 28, No. 1, pp. 37–50, 2000.
- [4] M. Beynon, D. Cosker, D. Marshall, *An expert system for multi-criteria decision making using Dempster-Shafer theory*, Expert Systems with Applications, Vol. 20, No. 4, pp. 357–367, 2001.
- [5] E.H. Forman, S.I. Gass, *The analytical hierarchy process: an exposition*, Operations Research, Vol. 49, No. 4 pp. 46–487, 2001.
- [6] R.D. Holder, *Some Comment on the Analytic Hierarchy Process*, Journal of the Operational Research Society, Vol. 41, No. 11, pp. 1073–1076, 1990.
- [7] F.A. Lootsma, *Scale sensitivity in the multiplicative AHP and SMART*, Journal of Multi-Criteria Decision Analysis, Vol. 2, pp. 87–110, 1993.
- [8] J.R. Miller, *Professional Decision-Making*, Praeger, 1970.
- [9] J. Perez, *Some comments on Saaty's AHP*, Management Science, Vol. 41, No. 6, pp. 1091– 1095, 1995.

- [10] T. L. Saaty, *Multicriteria Decision Making, The Analytic Hierarchy Process, Planning, Priority Setting, Resource Allocation*, RWS Publications, Pittsburgh, USA, 1988.
- [11] T. L. Saaty, *Decision-making with the AHP: Why is the principal eigenvector necessary*, European Journal of Operational Research, 145, pp. 85-91, 2003.
- [12] Octagon Mathematical Magazine (1993-2025)

Emeritus Professor,
Mathematics, Physics, and Natural Science Division,
University of New Mexico,
705 Gurley Ave., Gallup, NM 87301, USA